



Effects of long-term feeding of corn co-products on composition, tenderness, and shelf-life stability of beef strip steaks



J. R. Segers, R. L. Stewart Jr., T. D. Pringle, M. A. Froetschel, R.O. McKeith, B.K. Lowe, and A. M. Stelzleni

Department of Animal and Dairy Science, University of Georgia, Athens 30602

INTRODUCTION

- Recent volatility of feed prices and economic instability has forced beef cattle producers to re-evaluate nutritional programs
- Limited access to economical sources of corn and soybean meal has prevented the development of a feedlot industry in the SE
- Demand for locally-grown beef has increased interest in developing Southeastern beef cattle production systems utilizing corn co-products
- There is limited research on meat quality from cattle fed dried distillers grains plus solubles (DDGS) and corn gluten feed (CGF) from weaning to slaughter

OBJECTIVES:

- Evaluate performance and compositional development of steers fed corn co-products from weaning through slaughter
- Compare carcass characteristics, meat quality, and shelf-life of strip steaks from steers fed corn co-products

MATERIALS AND METHODS

- Angus-based beef steers (n=36) were fed at the UGA Wilkins Beef Cattle Research Unit for 100 d
- Individual intake was measured using a Calan® gate system
- Feedlot diets were formulated to be isonitrogenous so that protein supplement accounted for 25% of the diet
- Supplements included dry distillers grains with solubles (DDGS), corn gluten feed (CGF), and soybean meal with corn (SBM)
- After the 100-d feeding trial steers (n=36) were slaughtered at the UGA Meat Science Technology Center
- Carcass data were collected 24 h post-mortem by trained personnel
- At 48 h post-mortem the strip loin was removed and fabricated into 2.54-cm steaks
- Proximate analysis was used to determine moisture, protein, and lipid content
- Warner-Bratzler shear force was measured over 21-d aging period
- Steaks were also used to determine differences in shelf-life, and lipid oxidation over a 9-d aging period

CONCLUSIONS & IMPLICATIONS

- Feeding dried distillers grains or corn gluten feed yield similar meat quality characteristics to feeding soybean meal
- Dry distillers grains or corn gluten feed have no adverse effects on shelf-life or tenderness of strip steaks
- Corn co-products maybe utilized to develop a locally grown market for beef in the Southeast

RESULTS

Table 1. Carcass Yield Characteristics

Item ¹	Protein Supplement			SEM
	CGF	DDGS	SBM	
HCW, kg	354	350	341	7.33
DP,%	63.8	62.9	63.5	0.59
REA,cm ²	77.65	77.68	79.23	3.25
FT, cm	1.20	1.11	1.20	0.07
KPH, %	2.3	2.2	2.2	0.11
YG	3.10	3.11	3.05	0.18

¹DP=dressing percent, REA=ribeye area, FT=12th rib fat over the ribeye, YG=yield grade

Table 2. Carcass Quality Characteristics

Item	Protein Supplement			SEM
	CGF	DDGS	SBM	
Marbling ¹	494.2	432.5	457.5	23.44
Texture ²	1.42	1.75	1.50	0.17
Firmness ³	1.67	1.42	1.67	0.20
pH	5.57	5.64	5.59	0.05
Overall Maturity	A36	A31	A45	6.57

¹ 300=slight, 400=small, 500=modest,

² 1=very fine, 5=course

³ 1=very firm, 5=soft

Table 3. Proximate Analysis of Strip Steaks

Item	Protein Supplement			SEM
	CGF	DDGS	SBM	
Moisture, %	71.38	73.11	72.29	0.52
Protein, %	22.83	23.08	23.51	0.39
Lipid, %	5.31	3.83	4.22	0.54

Table 4. Meat Color Characteristics


Item	Protein Supplement			SEM
	CGF	DDGS	SBM	
Lean Color				
L*	43.73 ^a	41.21 ^b	40.67 ^b	0.78
a*	31.29	29.54	30.74	1.17
b*	12.68 ^a	10.96 ^b	11.02 ^b	0.38
Fat Color				
L*	73.08 ^{ab}	72.14 ^b	74.30 ^a	0.63
a*	10.32 ^a	9.47 ^{ab}	7.35 ^b	0.93
b*	16.50	13.37	14.46	1.38
Lean Color Score ¹				
Fat Color Score ²	1.83	1.61	1.75	0.17

^{a,b} Within a row means without a common superscript differ ($P < 0.05$)

¹ 1=extremely dark red, 8=extremely bright cherry red


² 1=white, 5=yellow

Figure 1. Warner-Bratzler Shear force




Shelf-life Trial

Figure 2. Subjective acceptability¹




¹ 8=extremely acceptable, 1=extremely unacceptable

Figure 3. Subjective color¹



¹ 8=light cherry red, 1=extremely dark red

Figure 4. Subjective discoloration¹



¹ 8=no discoloration, 1=complete discoloration

Figure 5. Lipid Oxidation




Figure 6. L*




Figure 7. a*




Figure 8. b*

